

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) **EP 0 637 203 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
**21.08.2002 Bulletin 2002/34**

(51) Int Cl.7: **A01N 45/00, A61K 31/56,**  
**A61K 38/21, A61K 45/00,**  
**C07J 1/00, A61K 31/565**

(21) Application number: **92915585.1**

(86) International application number:  
**PCT/US92/03076**

(22) Date of filing: **14.04.1992**

(87) International publication number:  
**WO 93/020696 (28.10.1993 Gazette 1993/26)**

### (54) REGULATION OF THE IMMUNE SYSTEM

REGULATION DES IMMUNSYSTEMS

REGULATION DU SYSTEME IMMUN

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU MC NL SE**

(74) Representative: **Harding, Charles Thomas et al**  
**D. Young & Co.**  
**21 New Fetter Lane**  
**London EC4A 1DA (GB)**

(43) Date of publication of application:  
**08.02.1995 Bulletin 1995/06**

(56) References cited:  
**US-A- 4 628 052 US-A- 4 898 694**  
**US-A- 4 956 355**

(60) Divisional application:  
**02075413.1 / 1 208 842**

(73) Proprietor: **LORIA, Roger**  
**Richmond, VA 23227 (US)**

• **ANN. N. Y. ACAD. SCI., vol. 650, 1992 pages**  
**363-366, R.M. LORIA ET AL. 'Mobilization of**  
**cutaneous immunity for systemic protection**  
**against infections.'**

(72) Inventor: **LORIA, Roger**  
**Richmond, VA 23227 (US)**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 637 203 B1**

**Description****Field of the Invention:**

[0001] The present invention provides an improved means for regulating the immune response, for ameliorating effects of stress, and for avoiding untoward effects of chemotherapy or exposure to irradiation by administration of androstenediol (AED) and androstenetriol (AET). The improved means of regulating immune response can be utilized in treating infectious diseases and immune diseases such as diabetes and chronic fatigue syndrome, both diseases now considered to be immune response related syndromes.

**Background of the Invention:**

[0002] In vertebrates the development of host protection against pathogens requires a selective host immune response that involves the mobilization of the humoral and/or cellular mediated immune responses. Several factors adversely affect the body's protective response capability by causing prolonged immuno-suppression or "down-regulation" of the immune system. It is, in reality, more appropriate to speak of "mal-regulation" or "deregulation" of the immune system than of "down-regulation" since the result is a failure to protect the body from assault. Immuno-suppression provides an opportunity for pathogens to grow in the host. It does not matter what causes the primary insult to immunity. The resulting inability to muster the appropriate immune response has the same effect. Among the many different causes of immuno-suppression are viral, bacterial, fungal, yeast and parasitic infections, chemotherapy, irradiation, severe stress, chronic fatigue syndrome, diabetes mellitus and some forms of steroid therapy.

[0003] It has long been known that patients receiving steroid hormones of adrenocortical origin at pharmacologically appropriate doses show increased incidence of infectious disease. A.S. Fauci, *immunolo.rev.* 65, 133-155 (1982); and J.E. Parillo and A.S. Fauci, *Annual Review of Pharmacology and Toxicology* 19, 179-201 (1979). Dehydroepiandrosterone, also known as 3- $\beta$ -hydroxyandrost-5-en-17-one or dehydroiso-androsterone (referred to hereinafter as DHEA), is a 17-ketosteroid which is quantitatively one of the major adrenocortical steroid hormones found in mammals. M.E. Windholz, *The Merck Index*, Ninth Edition (1976); K. Diem and C. Lentner, *Geigy Scientific Tables* (1975). (Although DHEA appears to serve as an intermediary in gonadal steroid synthesis, the primary physiological function of DHEA has not been fully understood. It has been known, however, that levels of this hormone begin to decline in the second decade of life, reaching 5% of the original level in the elderly.)

[0004] Clinically, DHEA has been used systemically and/or topically for treating patients suffering from psoriasis, gout, hyperlipemia, and it has been administered to post-coronary patients. W. Regelson et al., *New York Academy of Sciences* 518, 260-273 (1988). In mammals DHEA has been shown to have weight optimizing and anticarcinogenic effects.

[0005] DHEA has been used clinically in Europe in conjunction with estrogen as an agent to reverse menopausal symptoms and also has been used in the treatment of manic depression, schizophrenia, and Alzheimer's disease. DHEA has also been used clinically at 40 mg/kg/day in the treatment of advanced cancer and multiple sclerosis. (Regelson, *supra*). Mild androgenic effects, hirsutism, and increased libido were the side effects observed. These side effects can be overcome by monitoring the dose and/or by using analogues.

[0006] U.S. Patent 5,077,284 entitled "Use of Dehydroepiandrosterone to Improve Immune Response" describes the subcutaneous or oral administration of DHEA to improve the host's response to infections. U.S. Patent 4,978,532 describes use of patch technology to deliver DHEA.

[0007] It is now disclosed that DHEA is a precursor in a metabolic pathway which ultimately leads to more powerful agents that increase immune response in mammals. That is, DHEA acts as a biphasic compound: it acts as an immunomodulator when converted to androstenediol (5-androstene, 3 $\beta$ ,17 $\beta$  diol hereinafter referred to as  $\beta$ AED) or androstenetriol (5 androstene 3 $\beta$ ,7 $\beta$ ,17 $\beta$  triol hereinafter referred to as  $\beta$ AET), but *in vitro* has a certain lymphotoxic and suppressive effect on cell proliferation prior to its conversion to  $\beta$ AED and/or  $\beta$ AET. It is, therefore, now understood that administration of DHEA shows superior immunity enhancing properties as a result of its partial conversion to a more active metabolite.

[0008] An agent that would advance the protective regulation of the immune system without giving rise to undesirable side effects seen with DHEA administration would provide particularly advantageous improvement of host resistance against infection. Protective regulation of the immune system could then be effected using lower doses of the chemotherapeutic agent, and would provide more immediate response with a wider range of protection.

[0009] Roger M. Loria and David A. Padgett in N. Y. Acad. Sci., vol 650, pages 363-366 discuss androstenediol, a metabolite of dehydroepiandrosterone, and report that viral protection obtained by subcutaneous injection of androstenediol is quantitatively and qualitatively greater than that obtained with dehydroepiandrosterone.

**Description of the Invention**

[0010] According to the present invention there is provided 5-androstene-3 $\beta$ , 7 $\beta$ , 17 $\beta$ -triol ( $\beta$ AET) of a hydroxy-protected derivative thereof wherein the hydroxy groups are protected by hydrolysable protecting groups for use as an immune response modifier or regulator in man or other vertebrate species. The  $\beta$ AET may, for example, be present in admixture or in combination with a pharmaceutically or veterinarily acceptable carrier.

[0011] The present invention also provides the use of 5-androstene-3 $\beta$ , 7 $\alpha$ , 17 $\beta$ -triol ( $\alpha$ AET) for the manufacture of an immune response modifier or regulator in man or other vertebrate species. The  $\alpha$ AET may, for example, be used in admixture or in combination with a pharmaceutically or veterinarily acceptable carrier.

[0012] The present invention further provides a hydroxy-protected derivative of 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED) wherein the hydroxy groups are protected by hydroxy-protecting groups for use as an immune response modifier or regulator in man or other vertebrate species. The hydroxy-protected derivative of  $\beta$ AED may, for example, be present in admixture or in combination with a pharmaceutically or veterinarily acceptable carrier.

[0013] The hydroxy-protecting groups may, for example, be selected from C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl, phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl and -COR<sub>2</sub>, where R<sub>2</sub> is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl; phenyl or phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl; "alkyl" in such definitions including straight and branched chain alkyl, and wholly or partially cyclised alkyl groups, and "phenyl" including substituted phenyl groups containing up to 3 substituents selected from OH, C<sub>1</sub>-C<sub>4</sub> carboxy, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy and C<sub>2</sub>-C<sub>4</sub> alkenyl.

[0014] In the above mentioned mixtures or combinations according to the present invention the carrier may, for example, be a subcutaneously administrable carrier, a topically (including percutaneously) administrable carrier, or an orally administrable carrier. The subcutaneously administrable carrier may, for example, be a lipophilic carrier, and the topically administrable carrier may, for example, be a solution, cream, emulsion, suspension, jelly, powder, spray, patch or suppository.

[0015] The invention still further provides a method of preventing cell adhesion, which comprises contacting the cells with 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED), 5-androstene-3 $\beta$ , 7 $\beta$ , 17 $\beta$ -triol ( $\beta$ AET) or 5-androstene-3 $\beta$ , 7 $\alpha$ , 17 $\beta$ -triol ( $\alpha$ AET).

[0016] The present invention still further provides a method of enhancing lymphocyte proliferation *in vitro* which comprises incorporation into the growth medium 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED), or 5-androstene-3 $\beta$ , 7 $\beta$ , 17 $\beta$ -triol ( $\beta$ AET). The method may, for example, be applied to enhancing the proliferation of an autogenous lymphocyte population from a patient prior to readministration of the enhanced lymphocyte population to the patient.

[0017] The present invention still further provides a method for the substantially stereo-specific production of 5-androstene-3 $\beta$ , 7 $\beta$ , 17 $\beta$ -triol, which comprises the step of reducing 3 $\beta$ , 17 $\beta$ -diacetoxyandrost-5-ene-7-one with aluminium isopropoxide in isopropanol.

[0018] The present invention still further provides a method for the substantially stereo-specific production of 5-androstene-3 $\beta$ , 7 $\beta$ , 17 $\beta$ -triol which comprises the step of reducing 3 $\beta$ , 17 $\beta$ -diacetoxyandrost-5-ene-7-one with triisobutylaluminium in pentane to produce 3 $\beta$ , 17 $\beta$ -diacetoxy-7 $\beta$ -hydroxyandrost-5-ene, and subsequently removing the acetoxy groups.

[0019] The present invention still further provides, for example for use as an anti-viral agent, a composition comprising 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED) in a dosage of from 0.2 to 30 mg per day in a pharmaceutically acceptable carrier, the pharmaceutically acceptable carrier comprising a solid support. The support may, for example, comprise a patch or be a condom. Such composition may, for example, be used to contact a vaginal wall.

[0020] The present invention still further provides a capsule comprising 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED) in a dosage of from 0.2 to 30 mg per day in a pharmaceutically acceptable carrier.

[0021] The present invention still further provides the use of 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED) for the manufacture of a medicament for the prevention or reduction of damage by irradiation in man or other vertebrates species.


[0022] The present invention still further provides 5-androstene-3 $\beta$ , 7 $\alpha$ , 17 $\beta$ -diol ( $\alpha$ AET) for use as an active agent to prevent or reduce damage by irradiation in man or other vertebrate species.

[0023] The present invention provides compounds and compositions useful for enhancing the protective response of the immune system against infections. The medicinal compositions of the invention are also useful for treating other complications often accompanying immune suppression. The enhancement of the protective immune response may also be referred to herein as up-regulation or regulation of immune response.


[0024] The present invention provides means of synthesizing both  $\alpha$  and  $\beta$  stereoisomers of AET. Both isomers are present in the body. However, since the activity of the isomers varies, it is important that means of making the separate isomers are provided. It is now known that the  $\beta$ AET is the active agent in regulation of immune response. It is, however, probable that the  $\alpha$ AET, when in excess, is in equilibrium with  $\beta$ AET. However, because it is not known at this time what the response delay interval is, the  $\beta$ AET is the composition that is the isomer used therapeutically.

[0025] Using  $\beta$ AED and  $\beta$ AET and protected analogues avoids certain side effects resulting from use of DHEA. The use of  $\beta$ AED and  $\beta$ AET avoids many of the androgenic side effects that occur when the precursor DHEA is administered. With  $\beta$ AED and  $\beta$ AET it is possible to obtain rapid, controlled enhancement of immune response. Furthermore, as

shown by the information below in Table 1, the  $\beta$ AED is unexpectedly superior to DHEA for purposes of protecting against viral assault since the protection offered differs both quantitatively and qualitatively from protection effected by DHEA. The structures of  $\beta$ AED and the  $\alpha$  and  $\beta$  isomers of AET are given below:



[0026] In addition to the medicinal compositions, the invention encompasses analogues and derivatives of  $\beta$ AED and  $\beta$ AET. As indicated later in the exemplification of synthetic techniques, some of the substituted compounds are useful as starting materials or intermediates in making  $\beta$ AED and  $\beta$ AET.  $\beta$ AED is available from commercial sources and has been used as a raw material for production of other steroids. (See U.S. Patent 2,521,586.) The product used in testing was obtained from Sigma.  $\beta$ AED and  $\beta$ AET may also be substituted with protective groups which, on hydrolysis yield  $\beta$ AED or  $\beta$ AET. Hence, acylated and alkylated derivatives are useful as precursors to the  $\beta$ AED and  $\beta$ AET. Compounds, encompassed by the present invention, include those of the formula:



wherein  $R_1$  may be H, alkenyl of 2-8 carbons, alkyl of 1-8 carbons, phenylalkyl of 1-4 carbons, phenyl or  $COR_2$ , wherein  $R_2$  is H; alkyl of 1-8 carbons, alkenyl of 2-8 carbons, phenylalkyl wherein the alkyl has 1-4 carbons (including benzyl) or phenyl. Any phenyl moiety may have up to three substituents chosen from among hydroxy, carboxy of 1-4 carbons, halo, alkoxy of 1-4 carbons, alkyl of 1-4 carbons, or alkenyl of 2-4 carbons and wherein any alkyl may be a straight chain, branched chain, or the alkyl may be wholly or partially cyclized.

[0027] The acute virus infection used as a model in the examples was chosen because of its widespread effects and the severity of the model to show protection. The Coxsackievirus B4 is lethal to mice. In man, Coxsackieviruses cause such varied pathologies as upper respiratory infection, pharyngitis, hemorrhagic conjunctivitis, meningitis, exanthem, encephalitis, hepatitis, infantile diarrhea, paralysis, pericarditis, and myocarditis. It is now believed that viruses of this group also have a role in the onset of juvenile diabetes.

[0028] The use of  $\beta$ AED and  $\beta$ AET and protected analogues as taught herein provides high levels of protection to vertebrates, including humans, against morbidity arising from infections or exposure to immune suppressive influences. In clinical medicine, treatment with  $\beta$ AED and  $\beta$ AET can lower morbidity in patients exposed to pathogenic organisms. These agents can be effectively used prophylactically in patients known to be particularly susceptible to infection. Patients undergoing surgery or chemotherapy or patients suffering from burns, hypoplastic or aplastic anemias, or diabetes are such susceptible patients who would benefit from prophylactic administration of  $\beta$ AED and/or  $\beta$ AET. Also among the causes of immuno-suppression are viral, bacterial, fungal, yeast and parasitic infections, chemotherapy, irradiation, severe stress, chronic fatigue syndrome and some forms of steroid therapy. The compositions of the invention are particularly useful for treating patients suffering from infections caused by viruses that destroy the body's immune response, such as human immunodeficiency virus (HIV) and hepatitis.

[0029] Both  $\beta$ AED and  $\beta$ AET are most efficiently formulated using lipophilic carriers such as DMSO. For subcutane-

ous administration to animals used in the examples, the active agents were dissolved 1:1 DMSO/ethanol, then diluted for subcutaneous administration to the animals. When the compositions were administered by mouth, they were added to chow to provide a composition containing .4% AED. For application to the skin, the AED may, for example, be dissolved in carrier material containing DMSO and alcohol, then applied to a patch. In such instances, the AED in solution may be added to another carrier such as glycerol before application of the composition to the support material of the patch. For vaginal or rectal administration, the AED may be administered by suppository, enema, or by application of creams, etc. Compositions of the invention may be administered by any method that will result in contact of the active agent with tissue of ectodermal origin. Such methods include subcutaneous or intradermal injection or topical application. One means of topical application is the use of skin patches impregnated with the active agent. This means of delivery is advantageous since it is non-invasive and easily administered by relatively unskilled care providers. Compositions of the invention can also be used in veterinary medicine to prevent morbidity that occurs during stress of shipping. Administration of  $\beta$ AED and  $\beta$ AET can be effective as a means to prevent spread of infectious disease and introduction of infectious organisms into the foods for human consumption.  $\beta$ AED and  $\beta$ AET can be administered by subcutaneous injection, in food or drink, by patches applied to the skin, or by inhalation. A particular health concern is the spread of infection through eggs. Eggs are frequently infected during development in the hen. Compositions containing active agents of the invention may be added to the feed or water to prevent bacterial infection in the eggs.

**[0030]** When patches are used on animals or birds the skin should be exposed directly to the patch. When a patch is used, it may be necessary to pluck or shave the bird or animal to expose the skin.

**[0031]** Other preferred methods of administration include buccal, sublingual, nasal or endotracheal routes. Sprays may be useful for this purpose. For nasal administration, the active agent may be delivered as a powder that is snorted. Inclusion complexes such as cyclodextrin inclusion complexes are appropriate compositions for administration to the oral-pharyngeal and nasal mucosa.

**[0032]**  $\beta$ AED and  $\beta$ AET may also be given with vaccines to enhance immune response. These agents may be administered either in a composition containing the vaccine or may be given in a composition separate from the vaccine.

**[0033]** The compounds of the invention may also be administered to the intestinal mucosa by oral or rectal routes. Suppositories, solutions for use as retention enemas, and creams or jellies are appropriate carriers for use in rectal administration.

**[0034]** Compounds of the invention may be applied to the vaginal mucosa using creams, jellies, suppositories, or douching solutions. In order to enhance immune response at the site of exposure to infectious organisms, the compounds may be added to prophylactic vaginal preparations or may be used as lubricants on condoms.

**[0035]** Administration of compositions of the invention has proven to be highly effective as a means of protecting against encephalitis and meningitis. The compositions of the invention may be administered intrathecally either at the spinal level or into the cisterna magna.

**[0036]** Active agents of the invention may be administered via ocular route using compositions in the form of drops, creams, or gel solutions or suspensions adapted for ocular application.

**[0037]**  $\beta$ AED and  $\beta$ AET inhibit the adherence properties of body cells. For purposes of effecting the anti-adherence properties the active agents of the invention may be delivered directly to the epithelial tissue during surgery. An example of such use would involve the application of compositions containing the active agents of the invention to the omentum in conditions such as infection, endometritis and malignancies of the bowel and ovary wherein adherence of foreign cells or particles to normal cells of the peritoneal lining is a problem. Compositions of the invention could, for example, be administered as mists or sprays.

**[0038]** It has now been shown *in vivo* that DHEA is converted to the  $\beta$ AED and both isomers of AET. The  $\beta$ AED and  $\beta$ AET then act as regulators of the immune response. In the skin, the conversion of DHEA to AED and subsequently to AET appears to be one of the metabolic pathways of DHEA. The human skin has the enzymatic machinery to form 7 $\beta$ -OH DHEA and to cause 7 $\beta$  hydroxylation of AED to yield  $\beta$ AET. While the human adrenal cortex and liver can form only 7 $\alpha$ -hydroxylation of DHEA, but not 7 $\beta$ -hydroxylation. The following table indicates the metabolic pathways of DHEA.

Trivial or common name		systematic name
(a)	Dehydroepiandrosterone	(DHEA) 3 $\beta$ -Hydroxy-5-androsten-17-one
(b)	Dehydroepiandrosterone sulfate	(DHEAS) 3 $\beta$ -Hydroxy 5 androsten-17-one-sulfate
(c)	7 $\beta$ -OH Dehydroepiandrosterone--	-----
d)	Androstenediol	(AED) 5-androstene-3 $\beta$ ,17 $\beta$ ,diol
e)	Androstenediol	(AET) 5-androstene-3 $\beta$ ,7 $\beta$ ,17 $\beta$ ,triol
(f)	---	7 keto-5- androstene-3 $\beta$ ,17 $\beta$ , diol
(g)	---	5-androstene-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ ,triol

(continued)

Trivial or common name		systematic name
(h)	---	5-androstene-3 $\beta$ , 17 $\beta$ , diol-3-sulfate
(i)	Testosterone	17 $\beta$ -Hydroxy-4-androstene-3-one
(j)	Androstenedione	4-androstene-3,17 dione
(k)	Dihydrotestosterone	17 $\beta$ -Hydroxy-5 $\beta$ androstane-3-one
(l)	Androsterone	3 $\beta$ -Hydroxy-5 $\beta$ androstan-17-one

[0039] DHEA (a) is known to have immune enhancing activity. However, dehydroepiandrosterone sulphate, "(b)", has been found to have no enhancing effect on the immune system. Both 5 androstene-3 $\beta$ , 11 $\beta$ , 17 $\beta$ ,-triol and 5 androstene-3 $\beta$ , 16 $\beta$ , 17 $\beta$ ,-triol have immunosuppressive effects. Testosterone, "(i)", is known to have an effect on the host immune response, but this effect does not result in protection from lethal infections. The qualitative and quantitative effects of testosterone and the other sex hormones on the immune response are both of a different nature and scope.

#### Example 1:

[0040] Use of androstenediol (5 androstene-3 $\beta$ , 17 $\beta$ , diol,  $\beta$ AED) results in marked and significant resistance against viral and bacterial infection. Dose curve experiments were conducted in the following manner:  $\beta$ AED and DHEA were administered as a single depo dose of 8.3 mgs AED or 25 mgs. DHEA to SWRJ and C57BL/6J inbred mice. The mice were then challenged with varying amounts of Coxsackievirus B4 (CB4) to determine protective value of the active agents.  $\beta$ AED provided 50% protective dose against coxsackievirus B4, which was as much as 100 times greater than protection provided by DHEA. In addition to the difference in ED50, the degree of protection against virus mortality achieved with androstenediol was also greater than the one obtained with DHEA injection. The increased protection effected by  $\beta$ AED against virus induced mortality was statistically significantly different that the protection obtained by DHEA ( $P < 0.05$ ).

TABLE 1

DHEA AND AED IN SWRJ MICE MORTALITY PER GROUP*		
CB4 10 <sup>5</sup>	VIRUS ALONE 1/6	DHEA 0/6 AED 0/6
CB4 10 <sup>6</sup>	VIRUS ALONE 3/6	DHEA 0/6 AED 0/6
CB4 10 <sup>7</sup>	VIRUS ALONE 5/6	DHEA 1/6 AED 0/5
CB4 10 <sup>8</sup>	VIRUS ALONE 4/6	DHEA 1/6 AED 0/6

\*No deaths occurred in control groups.

AED versus control p-value = 0.0001.

DHEA versus control p-value = 0.0017.

DHEA versus AED p-value = 0.0588.

[0041] As seen from these results,  $\beta$ AED is markedly more efficient than the precursor DHEA in preventing mortality since an effective dose of  $\beta$ AED which is 1/3 or less the dose necessary to obtain an effect with DHEA is effective in achieving protection from mortality. A similar protection from virus mediated mortality was also observed in the inbred C57BL/6J(b) strain. The two inbred mouse strains, the SWRJ and the C57BL/6J, differ in their major histocompatibility haplotypes, which are q and b respectively. These results show that up-regulation of the immune response achieved with  $\beta$ AED in strains of different histocompatibility may be independent of the major histocompatibility genes on chromosome 17.

[0042] Recent reports show that the skin may have unique immune functions. J.W. Streilein and R.E. Tigelaar, *Photoimmunology*, Parrish et al. eds. (Plenum Publishing, New York, 1983) pp. 95-130. Indeed the skin is known to contain a population of cutaneous immune cells, which include the epidermal Langerhans cells and keratinocytes that produce an epidermal thymocyte-activating factor, similar to IL-1, in the murine system's Thy-1+ dendritic epidermal cell.

#### Example 2:

[0043] A composition containing 8 mg  $\beta$ AED was administered subcutaneously to outbred ICR mice who were then challenged with a lethal dose of *Streptococcus faecalis* strain X1515. OG1R Animals given  $\beta$ AED showed marked resistance to morbidity as evident from reduction in mortality from 57% in animals challenged with bacteria only, to 0% mortality in animals infected and treated with a single subcutaneous (SC) dose of  $\beta$ AED. Moreover,  $\beta$ AED above a

certain threshold dose mediates a considerable proliferation of lymphocyte cells in the spleen and thymus, but only in infected animals. Administration of 8 mg/animal  $\beta$ AED without exposure to the virus did not cause proliferation. Histopathological examination of the organs of inbred SWR/J mice infected with virus only and animals treated with  $\beta$ AED only, or  $\beta$ AED treated and virus infected revealed that  $\beta$ AED is effective in protecting from virus-induced myocardiopathy, and pancreopathy. Data presented in Table 2 below, shows that  $\beta$ AED protects the host ICR inbred strain (Inst. of Cancer Research strain now supplied by Holland Sprague Dowley Company) from *S. faecalis*, but at a dose which is 1/3 the dose of DHEA required for the same effect.

TABLE 2

BACTERIUM	DOSE			%
	AGENT	mg/ANIMAL	24 hr.	MORTALITY
<i>S. faecalis</i> .	none	0	4/7	57%
<i>S. faecalis</i>	DHEA	25	0/7	0
<i>S. faecalis</i>	AED	8.3	0/7	0

[0044] Protection was accomplished in an extremely acute infection where deaths occurred within 24 hours (4/7 deaths in non-treated groups versus 0/7 in the treated). Mice were challenged with a lethal dose of plasmids containing bacterium *S. faecalis* isolate X1515.0GIRF.

**Example 3:**

[0045] Wells containing monolayered urinary bladder tumor cells were covered with a 50  $\mu$ molar solution of either  $\beta$ AED or DHEA overnight before being used in the assay. The results are shown in Table 3 below.

TABLE 3

WELL	X1515.0GIRF DOSE	NO. ADHERED	% ADHERED
Media	$1.78 \times 10^6$	$1.77 \times 10^6$	99.4
DMSO	$1.78 \times 10^6$	$1.98 \times 10^6$	111.2
DHEA	$1.78 \times 10^6$	$1.01 \times 10^6$	56.7
AED	$1.78 \times 10^6$	$1.05 \times 10^6$	59.0

[0046] As can be seen from Table 3, both DHEA and  $\beta$ AED inhibit the adherence of the bacteria to human urinary bladder tumor cells. It is believed that  $\beta$ AED and AED have an effect on either membrane fluidity, or on a component which influences adhesion and/or penetration into the cell. Urinary Bladder Tumor cells (EJ6) were analyzed by flow cytometry using the fluorescent probe 1-4, trimethylammonio-phenyl-6 1,3,5 hexatriene (TMA-DPH) as a membrane probe to determine the effects of AED on membrane fluidity.

Result	
A.	Samples 1 and 2 represent CONTROL cells grown in standard tissue culture media.
B.	Samples 3 and 4 represent cells grown in the above control media with 50 $\mu$ M AED added.
	Sample A 1. 0.121256
	Sample A 2. 0.120548
	CONTROL AVERAGE 0.120902
	AED 50 $\mu$ M 3. 0.135151
	AED 50 $\mu$ M 4. 0.142453
	TEST AVERAGE 0.138802

[0047] The difference of 0.017899 between the control and  $\beta$ AED-treated cells is significant and demonstrates that  $\beta$ AED caused a significant change in cell membrane adhesion. It may be that an increase in cell membrane rigidity (decrease fluidity) influences adherence.



**Example 4:**

[0048] Cell culture media was inoculated with  $4 \times 10^7$  Hela cells. DHEA was added to a concentration 50  $\mu$ M and  $\beta$ AED was added to a concentration of 50  $\mu$ M. The cell culture exposed to DHEA showed a two-fold decrease in cell number. The cell culture exposed to the  $\beta$ AED, the control, and the culture exposed to the carrier alone showed a two fold increase or no change in the number of viable cells. These observations indicate  $\beta$ AED lacks deleterious effects of DHEA during the initial phase of cell response. Neither DHEA nor  $\beta$ AED had an effect on the number of virus infectious particles in vitro.

**Example 5:**

[0049] Mice were injected with 8 mg  $\beta$ AED or 25mg DHEA subcutaneously. The mice were then infected with Cox-sackievirus B4,  $10^5$  particles. The mice were sacrificed after 7 days. The spleen lymphocytes were removed and stimulated with Conavalin A, 5  $\mu$ g/ml in vitro. Stimulation was measured by  $^3$ H-thymidine incorporation method. The  $\beta$ AED had a profound effect on the proliferation of spleen lymphocytes with as much as 6.6 fold greater proliferation evidenced. The data also indicates that, during the initial phase following administration of the steroid, DHEA results in some suppression of the immune system.

[0050] The results described above indicate clearly the advantage of using  $\beta$ AED or  $\beta$ AET rather than DHEA to increase immune response to infections, since the initial immune suppression or toxicity that precedes the immune up-regulation that is seen in treatment with DHEA is avoided. Hence, therapy with  $\beta$ AED and  $\beta$ AET, rather than DHEA is preferred to improve immune response to infectious organisms.

[0051] The only change in structure between DHEA and AED is a reduction of the keto group at the 17 position to a  $\beta$  hydroxyl group. The AET which has the hydrogen at the 7 position replaced with a  $\beta$  hydroxyl group is produced by a metabolic pathway in the skin. In other words,  $\beta$ AET is a down-stream product of metabolic pathway from DHEA in the skin. The  $\beta$ AET appears to be the most effective compound for effecting immune up-regulation and protection from untoward effects of stress, chemotherapy, and irradiation.

**Preparatory method #1**

[0052] An important aspect of the invention is the preparation of sterio-specific  $\beta$ AET for use as a medicinal. The synthesis was accomplished using the 7-oxo- $3\beta$ , 17 $\beta$  acetoxo-androst-5-ene as a starting material.

SYNTHESIS OF  $3\beta$ , 7 $\beta$ , 17 $\beta$ -TRIHIDROXYANDROST-5-ENE(I) and  $3\beta$ , 7 $\alpha$ , 17 $\beta$ -TRIHIDROXYANDROST-5-ENE(II):

[0053] Chromic oxide oxidation of  $3\beta$ , 17 $\beta$ -diacetoxandrost-5-ene in glacial acetic acid gave  $3\beta$ , 17 $\beta$ -diacetoxandrost-5-en-7-one, (III), the intermediate to (I) or (II).

[0054] Aluminum isopropoxide reduction of (III) in isopropanol gave (I). lithium tri (sec-butyl) borohydride reduction of (III) in tetrahydrofuran yielded (II).

**PREPARATION OF  $3\beta$ , 17 $\beta$ -DIACETOXYANDROST-5-EN-7-ONE(III)**

[0055] 37.4 g(.1 mol)  $3\beta$ , 17 $\beta$ -diacetoxandrost-5-ene (Steraloids A7850) in 400 ml glacial acetic acid was reacted with 30.06 g (.3 mol) chromium(VI) oxide (Aldrich 23,265-3) dissolved in 20 ml H<sub>2</sub>O and 20 ml glacial acetic acid. The CrO<sub>3</sub> solution was added drop-wise to the  $3\beta$ , 17 $\beta$ -diacetoxandrost-5-ene solution while maintaining the temperature at 55°C for 4 hours. In order to decompose any unreacted CrO<sub>3</sub>, methanol was added to the reaction mixture followed by aqueous salt solution and ether. Evaporation of the ether yielded 7.8 g(20% yield) of crude III. Crystallization from 95% EtOH yielded (III) m.p. 214-215°C, DSC peak 191-224°C, max at 220°C Lit., <sup>1</sup> m.p. 218-219°C, Lit., <sup>2</sup> m.p. 224-225°C (from methanol).

Normal phase tlc: EtOAc-cyclohexane-EtOH (45:45:10), Rf=0.86. IR bands (cm<sup>-1</sup>: 1737, 1666 (Lit., <sup>2</sup> 1728, 1668). <sup>1</sup>H NMR (CDCl<sub>3</sub>), (d), ppm: 0.81(s, 3H), 1.25 (s, 3H), 2.02(s, 6H), 2.25(m, H at C-4), 2.5(m, H at C-8), 4.6(t, H at C-17), 4.7(m, H at C-3), 5.72(s, 1H); (Lit., <sup>2</sup> 0.80(s, 3H), 1.20(s, 3H), 2.03 (s, 6H), 4.62(m, 1H), 5.71(g, 1H).) Reverse phase lc/ms(fast atom bombardment detection) detected m/z 389(M+H)<sup>+</sup> ion in the major lc peak, (Lit., <sup>2</sup> m/z 388(M)). Tentative C-13 nmr, (d), ppm assignments (CDCl<sub>3</sub>) are:

Carbon 1	2	3	4	5	6	7	8	9
ppm 38.35	35.74	72.04	43.03	164.22	126.44	201.18	65.82	49.69
Carbon 10	11	12	13	14	15	16	17	18
ppm 37.76	27.26	35.95	44.72	44.95	25.83	27.55	81.90	12.05
Carbon 19	20	21	22	23				
ppm 20.67	171.14	21.24	170.23	21.17				

PREPARATION OF 3 $\beta$ ,7 $\beta$ ,17 $\beta$ -TRIHIDROXYANDROST-5-ENE(I)

[0056] 3 $\beta$ ,17 $\beta$ -diacetoxyandrost-5-en-7-one was subjected to reduction by aluminum isopropoxide in isopropanol to give 3 $\beta$ ,7 $\beta$ ,17 $\beta$ -trihydroxyandrost-5-ene.

PREPARATION OF 3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -TRIHIDROXYANDROST-5-ENE(II)

[0057] 5.1 ml(5.1 mmol) lithium tri(sec-butyl)borohydride (Aldrich L-Selectride) in tetrahydrofuran was rapidly added to 499 mg(1.28 mmol) of 3 $\beta$ ,17 $\beta$ -diacetoxyandrost-5-en-7-one in 15 ml of freshly distilled tetrahydrofuran under nitrogen while stirring for 1.5 hours at ice-bath temperature. 0.9 g KOH in 15 ml methanol was added, the reaction mixture refluxed for 0.5 hours, and then 37.5 ml of 10% NaCl solution was added. After cooling in freezer(-20°), crystals formed which were filtered to yield 123.6 mg (19%) (II), m.p. 239-45°C. Crystallization from methanol yielded (II), m.p. 249.5-253°C. <sup>1</sup>H nmr (CD(OD)<sub>3</sub>), (d),ppm: 0.75(s,3H),1.01(s,3H), 3.1(m,1H), 3.6(t,1H),3.7(d,1H),5.50(d,1H).

Tentative C-13 nmr, (d),ppm assignments (CD(OD)<sub>3</sub>) are:

Carbon 1	2	3	4	5	6	7	8
ppm 37.49	32.12	72.01	42.91	146.75	124.88	65.46	39.09
Carbon 9	10	11	12	13	14	15	
ppm 43.58	38.55	21.49	30.65	93.65	45.35	38.13	

Carbon	16	17	18	19
ppm	42.57	82.30	11.57	19.53

## DISCUSSION

[0058] Stereochemistry was assigned to 3 $\beta$ ,7 $\beta$ ,17 $\beta$ -trihydroxyandrost-5-ene(I) and 3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -trihydroxyandrost-5-ene(II) by comparison with cholest-5-ene-3 $\beta$ ,7 $\beta$ -diol and cholest-5-ene-3 $\beta$ ,7 $\alpha$ -diol proton nmr(Lit.3). L-Selectride reduction of (III) to produce (II) was carried out using the same reaction conditions given by Morisaki for the preparation of 7 $\alpha$ -hydroxycholesterol<sup>4</sup>. Carbon-13 resonances for stereoisomers (I) and (II) are shown. (Multiplicities (ppm))

ISOMER(II)		ISOMER(I)	
1	37.19	T	37.77
2	32.12	T	32.30
3	72.01	D	72.12
4	42.91	T	41.24
5	146.75	*	144.10
6	121.88	D	127.43
7	65.46	D	74.00
8	39.09	D	38.29

(continued)

ISOMER(II)		ISOMER(I)	
9	43.58	D	50.08
10	38.55	*	37.73
11	21.49	T	21.90
12	30.65	T	30.86
13	43.65	*	44.27
14	45.35	D	52.31
15	38.31	T	26.60
16	42.57	T	42.57
17	82.30	D	82.30
18	11.57	Q	11.57
19	19.53	Q	19.53

Q=quartet, T=triplet, D=doublet and \*= quaternary C

(ISOMER II, 7-OH axial) (ISOMER I 7-OH equatorial)

## REFERENCES

## [0059]

1. Adolf Butenandt et al. Ber. 71B, 1316-22(1938).
2. Anthony J. Pearson et al., J. Chem. Soc. Perkin Trans. I, 267-273(1985).
3. Leland L. Smith et al., J.Org.Chem., Vol.38, No.1, 119-123 (1973).
4. Masuo Morisaki et al., Chem. Pharm. Bull. 35(5)1847-1852, (1987)

Preparatory method #2

[0060] A second method of preparing the  $\alpha$  and  $\beta$  isomers of AET has been developed. The process uses, as a starting material,  $3\beta$ ,  $17\beta$  diacetoxyandrost-5-ene, a commercially available reagent (Stereloids A7850). This compound is oxidized with chromium hexacarbonyl ( $\text{Cr}(\text{Co})_6$ ) to form  $3\beta$ ,  $17\beta$ -diacetoxyandrost-5-en-7-one as described by Pearson (Pearson, et al., J. Chem Soc. Perkin. Trans. 1, 1985, 267.) The enone formed was then reduced with triisobutylaluminum (TIBA) to give the acetylated  $7\alpha$ -hydroxy (3a) or  $7\beta$ -hydroxy (3b) product depending on the solvent used.

PREPARATION OF  $3\beta$ ,  $7\alpha$ ,  $17\beta$ -TRIHIDROXYANDOST-5-ENE.

[0061] A solution of  $3\beta$ ,  $17\beta$ -diacetoxyandrost-5-en-7-one (0.4991 g., 1.285 mmol) and tetrahydrofuran (20 mL, dried over  $\text{MgSO}_4$ ) were mixed under a nitrogen atmosphere. TIBA (2 mL, 2mmol, in 1M toluene) was added dropwise by syringe. The solution was stirred at room temperature for about 7 hours. The reaction was terminated by the addition of ethyl acetate (1.4mL), the methanol (10mL), and finally, by addition of 10 mL of 50% acetic acid. This solution was added to 100 mL water and extracted with ethyl acetate (3 times with 50 mL.) The organic layers were combined and then washed with saturated sodium bicarbonate solution (50 mL), saturated sodium chloride solution (2 times with 50 mL) and water (50 mL). The organic layer was then dried over magnesium sulfate and the solvent removed by rotary evaporation to yield >96% crude product.  $^1\text{H}$  NMR indicated that the crude product contained  $3\beta$ ,  $17\beta$ -diacetoxyandrost-5-en-7-one (starting material), 65% and  $3\beta$ ,  $17\beta$ -diacetoxy- $7\alpha$ -hydroxyandrost-5-ene, 35%.

PREPARATION OF  $3\beta$ ,  $7\beta$ ,  $17\beta$ -TRIHIDROXYANDROST-5-ENE

[0062] A solution of  $3\beta$ ,  $17\beta$  diacetoxyandrost-5-en-7-one (0.9581 g, 2.466 mmol) and pentane (30 mL), dried over  $\text{MgSO}_4$ , were mixed under a nitrogen atmosphere. TIBA (9.5 mL, 9.5 mmol, 1M in toluene) was then added dropwise by syringe. The solution was stirred at room temperature for about 1 hour. The reaction was terminated by the addition of diluted hydrochloric acid (approximately 5mL). This solution was added to water (100 mL) and extracted with ethyl acetate (3 times with 50 mL). The organic layers were combined and then washed several times with saturated sodium

bicarbonate solution (50 ml), saturated sodium chloride solution (two times with 50 mL,) and water (50 mL). The organic layer was dried over magnesium sulfate and the solvent removed by rotary evaporation to yield 86% crude product. <sup>1</sup>H NMR indicated that the crude product contained 3 $\beta$ ,17 $\beta$ , -diacetoxy-7- $\beta$ -hydroxyandrost-5-en-7-one (86%) and 3 $\beta$ ,17 $\beta$ -diacetoxy-7- $\alpha$ -hydroxyandrost-5-ene.

[0063] In both of the above instances, the final products were reduced in accord with preparatory method #1.

[0064] The compositions of the invention containing  $\beta$ AED and  $\beta$ AET, are useful in the treatment of any condition wherein a condition related to immune response is present including diabetes mellitus, chronic fatigue syndrome, stress, chemotherapy, and exposure to irradiation. One of the more depressing results of many of these conditions that involve aberrations of the immune response includes alopecia. The compounds of the invention are effective for treating the alopecia resulting from such aberrations of immune response, including effects of chemotherapy and irradiation. In alopecia, the  $\alpha$ AET should be of considerable benefit for use in treatment. The compositions of both  $\beta$ AED and  $\beta$ AET are useful in overcoming other untoward effects of immune suppression arising in these conditions.

#### Example 6:

[0065] SWR/J inbred mice were injected with single depo doses of 0.5, 2.0, 4.0 and 8.0 mg/animal of  $\beta$ AED, then challenged with 10<sup>7</sup> PFU Coxsackievirus B4. The results are shown below:

AED dose, mg/animal	% cumulative survival
0	17
0.5	83
2	100
4	100
8	100

[0066] Based on these results, the theoretical dose (extrapolated) of  $\beta$ AED necessary to achieve 50% protection from infection with 10<sup>7</sup> PFU of CB4 is 0.25 mg/animal.

[0067] The example demonstrates that  $\beta$ AED is 50 to 100 times more potent than DHEA for protection from Coxsackievirus B4.

#### Example 7:

[0068] In an attempt to evaluate effect of  $\beta$ AED on pathologies of the heart and pancreas in infected mammals, three groups of inbred SWR/J mice were compared. The groups studied included animals infected with virus only, animals treated with  $\beta$ AED only, and animals infected with virus and protected with  $\beta$ AED. The comparison revealed the following:

	Heart	Pancreas
AED	unremarkable	unremarkable
VIRUS	focal areas of multiple necrosis with substantial myocardial calcification.	severe necrosis (Pancreopathy)
AED PLUS virus	no evidence of induced myocardiopathy	no evidence of virus induced pancreopathy

[0069] These results demonstrate that  $\beta$ AED given as a single depo dose at 8.0 mg/mouse protected the heart tissue from virus induced myocardiopathy, and also protected the pancreas from virus induced necrosis of this organ. These results show that  $\beta$ AED can be used effectively in the protection from virus-induced cardiovascular and pancreatic disease, in particular myocardiopathies and pancreopathies. Previously there were no effective drugs to protect these organs from virus induced damage.

#### Example 8:

[0070] Effectiveness of  $\beta$ AED,  $\alpha$ AET, and  $\beta$ AET were compared by challenging mice with Coxsackievirus B4, 5.0 x 10<sup>7</sup> pfu/animal. The following data represents total cumulative mortality (n/6) at indicated days post infection with virus.

Treatment Group	Cumulative mortality							
	(Days Post Infection)							
	2	4	6	8	10	12	14	16
Virus alone	0	1	6	*	*	*	*	*
$\beta$ AED .5 mg/virus	0	0	1	1	6	*	*	*
$\alpha$ AET .5 mg/virus	0	1	4	4	4	4	4	4
$\beta$ AET .5 mg/virus	0	0	1	1	2	2	4	4

[0071] While the  $\alpha$ AET appeared to protect two animals from the virus, the initial mortality would indicate that at least in early stages, there is little or no protection from the  $\alpha$ AET. Other studies indicate that there is far less protection with  $\alpha$ AET than with either  $\beta$ AED or  $\beta$ AET. Clearly, the  $\beta$ AET is the most effective agent for both early and longer term protection. It appears that there is a limited protection attributable to  $\alpha$ AET. Therefore, this isomer would not be as beneficial as  $\beta$ AET for effecting enhancement of immune response.

[0072] The use of  $\beta$ AED or  $\beta$ AET as active agents to provide regulation of the immune system makes it possible to effectively regulate the host immune response against viral, bacterial and other infections. In the case of virus-induced heart or pancreatic infection where no other antiviral chemotherapeutic modality exists  $\beta$ AED and  $\beta$ AET have value as prophylactic protective agents. The protective value of  $\beta$ AED and  $\beta$ AET is particularly important to patients undergoing surgical procedures or suffering injuries where resistant strains of organisms such as pseudomonas present a serious threat. Examples of such patients are those undergoing bowel surgery or suffering from gunshot wounds of the abdomen. Patients with history of conditions such as rheumatic fever would also benefit from prophylactic use of  $\beta$ AED and  $\beta$ AET. The mode of administration in a particular case will depend on the infectious agent against which protection is sought and the target tissue of the organism. While the administration subcutaneously as a depo is effective against systemic infections as shown by the data presented above, when the compositions are given to assist the body in meeting an infection in a particular tissue, it may be advantageous to administer the active agents to the tissues most affected.

[0073] The carrier used in a given instance will depend on the mode of administration. Both AED and AET are lipophilic compounds. They are more soluble than DHEA in water. Solvents for lipophilic steroids are known in the art and would be used as carriers for these compounds. Examples of such carriers are glycols such as polypropylene glycol, polyethylene glycol, and cyclodextrins, especially the intrinsically amorphous cyclodextrins. Other vehicles that should be considered include fatty acid esters of polyoxyethylene sorbitan (Tweens) or sorbitan (Spans) to prepare oil-in-water emulsions.

#### Example 9:

[0074] Capsules of a formulation of AED for oral administration is prepared by containing 15 mg.  $\beta$ AED, 150 mg. starch, and 5 mg. magnesium stearate. The capsules are administered daily or twice a day to achieve a daily dosage of 15 mg. per day.

[0075] In the laboratory,  $\beta$ AED was added to the chow of the animals at a rate of .4% of the diet. When the animals who had been fed the diet containing  $\beta$ AED were exposed to Cocksackievirus B4 by injection in accord with the teachings of Example 1, the animals who had been fed the  $\beta$ AED survived, while control animals all died.

#### Example 10:

[0076] A preparation for application to the skin or mucosa may be prepared in the following manner:

Ingredient	%w/w
AED	0.5%
glyceryl monostearate	3.0%
propylene glycol	13.0%
Petrolatum	83.5%

[0077] When  $\beta$ AED or  $\beta$ AET or their analogues are administered orally, the active agents may be utilized more efficiently if the active agents are protected from destruction and absorption in the upper gastro-intestinal tract. The active agents are most effective when the period of exposure to the mucosa of the intestinal tract is increased. Hence

use of capsules containing the active agents in formulations that effect slow release in the intestine are appropriate for treatment of intestinal disorders such as Crohn's disease and colitis. Use of retention enemas for treatment of inflammation of the large bowel is also appropriate.

A formulation for administration as a retention enema may be formulated in the following manner:

**Example 11:**

**[0078]**

Ingredient	w/w %
$\beta$ AET	4%
Propylene glycol	96%

**[0079]** When the active agent is administered to the mucosa of the oral cavity, it may be administered as a buccal tablet or spray for use in the oral-pharyngeal cavity and the nasal cavities.

**[0080]** The compositions could also be administered to the bronchial tree via inhalation. This means of administration would be particularly useful in treating patients with lung infections or in treating other lung conditions such as black lung disease or emphysema that often are complicated by opportunistic infections. The compositions could be given by aerosol into the trachea or administered in- mist along with other agents used in respiration therapy.

**[0081]** The administration of the  $\beta$ AED and  $\beta$ AET to the skin can be accomplished using patches wherein a support is impregnated with the active agent or using implants that provide slow release of the active agents.

**[0082]** A patch for the administration of AET or AED can be formulated as adhesive patches containing the drug. For example, the patch may be a discoid in which a pressure-sensitive silicone adhesive matrix containing the active agent may be covered with a non-permeable backing. The discoid may either contain the active agent in the adhesive or may have attached thereto a support made of material such as polyurethane foam or gauze that will hold the active agent. Before use, the material containing the active agent would be covered to protect the patch.

**Example 12:**

**[0083]** A patch composed of trilaminate of an adhesive matrix sandwiched between a non-permeable backing and a protective covering layer is prepared in the following manner:

**[0084]** To a pressure-sensitive silicone adhesive composition BIOPSA™ Q7-2920 (Dow Corning Corp., Midland, Michigan, U.S.A.) in cyclohexane (50% w/v) is added sufficient  $\beta$ AED to provide a .5%  $\beta$ AED composition. The adhesive is applied to a polyester film to provide about 2 mg of active agent per cm<sup>2</sup>. The film containing the adhesive is then made into patches of 10 cm<sup>2</sup>. The patches would be covered with a protective layer to be removed before application of the patch. Patches may be prepared containing permeation enhancers such as cyclodextrin, butylated hydroxy-anisole, or butylated hydroxytoluene. However, it should be remembered that the active agents of this invention are effective on application to the epidermal tissue. When the patches are to be applied to thin or abraded skin, there is little need to add a permeation enhancer.

**[0085]** Compositions of the invention can be administered as a prophylactic during radiation therapy or chemotherapy or after exposure to irradiation whether the exposure occurs as a result of environmental accident or therapy. Other instances when use of these immune up-regulators would be appropriate is in treatment of burns, hypoplastic and aplastic anemias, diabetes, and in the elderly during epidemics. Their use is also beneficial in preventing or mitigating effects of exposure to dangerous infectious organisms, as was demonstrated by the data related to cardiopathies and pancreopathies. Such use is particularly indicated in populations exposed to organisms that target the immune system, such as HIV infections. In certain instances the compositions taught herein can also be used as immune modulators in the production of blocking antibodies to counteract hypersensitivity reactions.

**[0086]** As indicated previously, patients scheduled to undergo bowel surgery or other "dirty" surgical procedures could receive a dose of  $\beta$ AED or  $\beta$ AET prophylactically. Use of the compositions as taught herein before invasive dental procedures or oral surgery should be considered.

**[0087]** As indicated in the tables and previous discussion, the compositions of the invention can be used to prevent adhesion of bacteria to the tissues. The prevention of cell to cell adhesion resulting from administration of the compositions also has applications for prevention of thrombosis. Compositions of the invention can be administered as a slow drip into blood vessels when prevention of formation of a thrombus is necessary. An example of such use would be a drip into an artery following thrombolectomy or for prevention of cerebral thrombosis. Instillation into the bladder could also be beneficial for prevention or treatment of urinary infection. The administration of the  $\beta$ AED or  $\beta$ AET can be employed as a means of preventing formation of infectious foci.

[0088]  $\beta$ AED and  $\beta$ AET are both effective in overcoming effects of ultraviolet deprivation. Hence, administration of the compositions to overcome the effects of depression related to light deprivation (usually called seasonal adaptive disorder) is appropriate.

[0089]  $\beta$ AED and  $\beta$ AET may be used as adjuncts in vaccination to increase response to an immunogen. Not only will these agents increase response to the vaccine, they will also increase ability to protect against disease before the body has responded with increase in specific antibodies. Such use is particularly appropriate in instances where inhibition of immune response can be a complicating factor as is the case in patients suffering from, for example, malignancies, AIDS, or environmental factors such as exposure to pesticides. It is, of course, understood that use as adjunct to vaccination would be appropriate in vertebrates other than man, including vaccination of pets, dairy animals, meat-producing animals, fish, and chickens.

[0090] Chickens are particularly prone to develop infectious diseases when living in confined conditions. Coccidiosis, Salmonella infections, viral infections, including those giving rise to malignancies such as leukemia and sarcoma (caused by a retrovirus) are particularly common among chickens grown under modern commercial conditions. The active agents of the invention may be given by any means that results in contact of the agent with tissue of ectodermal origin.

[0091] The effect of Coxsackievirus on humans has been noted previously. The value of avoiding such effects, especially in children, is clear. Effects of other viruses such as chickenpox-herpes zoster is considered an important cause of debilitating illness in the elderly. Furthermore, chickenpox in susceptible adults often causes severe illness. In children, chicken pox can cause death when the child is subjected to immuno suppressive therapy or is genetically immune deficient.  $\beta$ AED and  $\beta$ AET are useful for prophylactic treatment of susceptible persons who have been exposed to infection. Finally, the protection of the fetus and newborn from HSV infection is a very important application of the invention.  $\beta$ AED and  $\beta$ AET can be administered during the third trimester to HSV-infected women as a means of protecting the newborn.

[0092] In vitro, these compounds can be used in commercial setting to induce lymphocyte proliferation. The use of  $\beta$ AED and  $\beta$ AET would increase yield of products of such proliferation in tissue culture. In the clinical setting,  $\beta$ AED and  $\beta$ AET can be given to effectively enhance patients' ability to combat infections. Patient lymphocytes may be withdrawn, reproduced in vitro in media containing  $\beta$ AED or  $\beta$ AET to increase proliferation of lymphocytes, and the lymphocytes so primed for response could then be reintroduced into the patient. In cases such as malignancy or other cellular disease, the malfunctioning cells can be inactivated by known means before proliferation in growth media.

[0093] The compositions of the invention may also be used prophylactically to protect animals from the consequences of infection by pathogenic organisms. It is known that under the stress of shipment to market animals often become susceptible to infections that are not ordinarily serious, but that can cause the animals to lose much weight en route to the packing house. Such loss may be avoided by administration of  $\beta$ AED and  $\beta$ AET and analogues disclosed herein. The active agents can be given by patch, injection, or in feed. Because the active agents are most effective when the period of exposure to the tissue of ectodermal origin is extended, when the active agents are administered through the GI tract, compositions should be modified to extend the period of exposure of the active agent to the intestinal mucosa and to protect the agents from destruction in the upper GI tract. Hence, use of capsules that effect slow release in the intestine is appropriate. The capsules may be placed in baits for administration to animals. To treat infections of the large bowel, the active agents may be given by retention enema.

[0094]  $\beta$ AED and  $\beta$ AET may be administered to the mucosa of oral, pharyngeal, and nasal cavity by tablet, a lozenge, by administration as a spray for use in the oral-pharyngeal cavity, or as a nasal spray.

[0095] Administration to the skin may be accomplished using patches wherein a support to be applied to the skin is impregnated with the active agent. If the host is a mammal or bird, it may be necessary to shave or pluck the region to which the patch is applied.

[0096] A preferred method of administration is by subcutaneous injection as a depo. The method is particularly appropriate for administration of the active agents to mammals, since subcutaneous injection is easily performed and the effect is relatively long lasting.

[0097]  $\beta$ AED and  $\beta$ AET are already present in the body as natural components. They do not pose a serious toxic problem at levels known to be safe; they appear to be chemically quite stable.

[0098] The dosages used will depend on the size and condition of the host. Test data indicated in this application was obtained in small animals. In larger adult mammals daily dosage of .2 to 30 mg/da. of AED is a preferred dosage. For AET the preferred dosage is usually in the range of .001 to 20 mg/da, with .001 to 1 mg/da. being the more preferred dosage. However, the dosage will vary depending on the route of administration. Subcutaneous, inhalation and intrathecal administration are methods that would require lower dosages of the active agents.

[0099] It is, of course, understood that analogues of  $\beta$ AED and  $\beta$ AET having protective groups can be administered to the host as a means of delivering  $\beta$ AED or  $\beta$ AET to target tissues. Acylation is a preferred method of protecting the compounds. Acylated compounds wherein  $R_1$  is  $COR_2$  are also appropriate compounds for use as starting material from which to make  $\beta$ AED and  $\beta$ AET.

[0100] The active agents,  $\beta$ AED and  $\beta$ AET, can be given in conjunction with other active agents which may be given simultaneously or may be incorporated in compositions containing  $\beta$ AED or  $\beta$ AET.  $\beta$ AED and  $\beta$ AET can be given with anti-infective agents such as antibiotics, antiviral agents, antifungals, antiparasitic agents to potentiate the activity of these drugs by up-regulating protective immune response. Antiviral agents include, for example, Dideoxyinosine, AZT, acyclovir, etc. Other active agents that may be combined with the AED and AET include antiallergic medications such as epinephrine.

[0101] Finally, medicinal compositions containing  $\beta$ AED and  $\beta$ AET are particularly valuable for use in combating viral infections in patients who have suffered from infections exacerbated by immuno-suppressive therapy. One of the major complications in patients with tissue transplants is the opportunistic infection with viruses that ordinarily do not cause serious disease symptoms. Use of the compositions of the invention, which result in rapid protective regulation of the immune response, allows the medical team to place the patient on "see-saw" therapy to avoid transplant rejection while regulating the immune response to avoid overwhelming infection.

## Claims

1. 5-androstene-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET) or a hydroxy-protected derivative thereof wherein the hydroxy groups are protected by hydrolysable protecting groups for use as an immune response modifier or regulator in man or other vertebrate species.
2. A product according to claim 1, wherein the hydroxy-protecting groups are selected from C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl, phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl and -COR<sub>2</sub>, where R<sub>2</sub> is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl or phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl; "alkyl" in such definitions including straight and branched chain alkyl, and wholly or partially cyclised alkyl groups, and "phenyl" including substituted phenyl groups containing up to 3 substituents selected from OH, C<sub>1</sub>-C<sub>4</sub> carboxy, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy and C<sub>2</sub>-C<sub>4</sub> alkenyl.
3. An admixture or a combination of a product according to claim 1 or 2 with a pharmaceutically or veterinarily acceptable carrier for use as an immune response modifier or regulator in man or other vertebrate species.
4. An admixture or a combination according to claim 3, wherein the carrier is a subcutaneously administrable carrier.
5. An admixture or a combination according to claim 4, wherein the subcutaneously administrable carrier is a lipophilic carrier.
6. An admixture or a combination according to claim 3, wherein the carrier is a topically (including percutaneously) administrable carrier.
7. An admixture or a combination according to claim 6, wherein the topically administrable carrier is a solution, cream, emulsion, suspension, jelly, powder, spray, patch or suppository.
8. An admixture or a combination according to claim 3, wherein the carrier is an orally administrable carrier.
9. Use of 5-androstene-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET) for the manufacture of an immune response modifier or regulator in man or other vertebrate species.
10. Use according to claim 9, wherein the hydroxy-protecting groups are selected from C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl, phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl and -COR<sub>2</sub>, where R<sub>2</sub> is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl or phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl; "alkyl" in such definitions including straight and branched chain alkyl, and wholly or partially cyclised alkyl groups, and "phenyl" including substituted phenyl groups containing up to 3 substituents selected from OH, C<sub>1</sub>-C<sub>4</sub> carboxy, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy and C<sub>2</sub>-C<sub>4</sub> alkenyl.
11. Use according to claim 9 or 10 wherein the immune response modifier or regulator comprises  $\alpha$ AET in admixture or in combination with a pharmaceutically or veterinarily acceptable carrier.
12. Use according to claim 11, wherein the carrier is a subcutaneously administrable carrier.
13. Use according to claim 12, wherein the subcutaneously administrable carrier is a lipophilic carrier.



14. Use according to claim 11, wherein the carrier is a topically (including percutaneously) administrable carrier.
15. Use according to claim 14, wherein the topically administrable carrier is a solution, cream, emulsion, suspension, jelly, powder, spray, patch or suppository.
- 5 16. Use according to claim 11, wherein the carrier is an orally administrable carrier.
- 10 17. A hydroxy-protected derivative of 5-androstene-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) wherein the hydroxy groups are protected by hydroxy-protecting groups for use as an immune response modifier or regulator in man or other vertebrate species.
18. A hydroxy-protected derivative of  $\beta$ AED according to claim 17 wherein the hydroxy-protecting groups are selected from C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl, phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl and -COR<sub>2</sub>, where R<sub>2</sub> is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, phenyl or phenyl (C<sub>1</sub>-C<sub>4</sub>) alkyl; "alkyl" in such definitions including straight and branched chain alkyl, and wholly or partially cyclised alkyl groups, and "phenyl" including substituted phenyl groups containing up to 3 substituents selected from OH, C<sub>1</sub>-C<sub>4</sub> carboxy, halo, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkoxy and C<sub>2</sub>-C<sub>4</sub> alkenyl.
- 15 19. An admixture or a combination of a hydroxy-protected derivative of  $\beta$ AED according to claim 17 or 18 with a pharmaceutically or veterinarily acceptable carrier for use as an immune response modifier or regulator in man or other vertebrate species.
- 20 20. An admixture or a combination according to claim 19, wherein the carrier is a subcutaneously administrable carrier.
21. An admixture or a combination according to claim 20, wherein the subcutaneously administrable carrier is a lipophilic carrier.
- 25 22. An admixture or a combination according to claim 19, wherein the carrier is a topically (including percutaneously) administrable carrier.
23. An admixture or a combination according to claim 22, wherein the topically administrable carrier is a solution, cream, emulsion, suspension, jelly, powder, spray, patch or suppository.
- 30 24. An admixture or a combination according to claim 19, wherein the carrier is an orally administrable carrier.
25. A method of preventing cell adhesion, which comprises contacting the cells with 5-androstene-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED), 5-androstene-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET) or 5-androstene-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET).
- 35 26. A method of enhancing lymphocyte proliferation *in vitro* which comprises incorporating into the growth medium 5-androstene-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED), or 5-androstene-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET).
- 40 27. A method according to claim 26, as applied to enhancing the proliferation of an autogenous lymphocyte population from a patient prior to readministration of the enhanced lymphocyte population to the patient.
28. A method for the substantially stereo-specific production of 5-androstene-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol, which comprises the step of reducing 3 $\beta$ ,17 $\beta$ -diacetoxyandrost-5-ene-7-one with aluminium isopropoxide in isopropanol.
- 45 29. A method for the substantially stereo-specific production of 5-androstene-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol which comprises the step of reducing 3 $\beta$ ,17 $\beta$ -diacetoxyandrost-5-ene-7-one with tri-isobutylaluminium in pentane to produce 3 $\beta$ ,17 $\beta$ -diacetoxy-7 $\beta$ -hydroxyandrost-5-ene, and subsequently removing the acetoxy groups.
- 50 30. A composition comprising 5-androstene-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) in a dosage of from 0.2 to 30 mg per day in a pharmaceutically acceptable carrier, the pharmaceutically acceptable carrier comprising a solid support.
31. A composition according to claim 30 wherein the support comprises a patch.
- 55 32. A composition according to claim 30 wherein the said support is a condom.
33. Use of composition according to any one of claim 30 to 32 to contact a vaginal wall.

34. A composition according to any one of claims 30 to 32 for use as an anti-viral agent.
35. A capsule comprising 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED) in a dosage of from 0.2 to 30mg per day in a pharmaceutically acceptable carrier.
36. Use of 5-androstene-3 $\beta$ , 17 $\beta$ -diol ( $\beta$ AED) for the manufacture of a medicament for the prevention or reduction of damage by irradiation in man or other vertebrates species.
37. 5-androstene-3 $\beta$ , 7 $\alpha$ , 17 $\beta$ -triol ( $\alpha$ AET) for use as an active agent to prevent or reduce damage by irradiation in man or other vertebrates species.

#### Patentansprüche

1. 5-Androsten-3 $\beta$ , 7 $\beta$ , 17 $\beta$ -triol ( $\beta$ AET) oder ein mit Hydroxylgruppenschutz versehenes Derivat davon, wobei die Hydroxylgruppen durch hydrolysierbare Schutzgruppen geschützt sind, zur Verwendung als ein Immunantwortmodifikator oder -regulator im Menschen oder anderen Wirbeltierspezies.
2. Produkt nach Anspruch 1, wobei die die Hydroxylgruppen schützenden Gruppen unter C<sub>1</sub>-C<sub>8</sub>-Alkyl, C<sub>2</sub>-C<sub>8</sub>-Alkenyl, Phenyl, Phenyl-(C<sub>1</sub>-C<sub>4</sub>)-Alkyl und -COR<sub>2</sub> ausgewählt sind, wobei R<sub>2</sub> H, C<sub>1</sub>-C<sub>8</sub>-Alkyl, C<sub>2</sub>-C<sub>8</sub>-Alkenyl, Phenyl oder Phenyl-(C<sub>1</sub>-C<sub>4</sub>)-Alkyl ist und wobei "Alkyl" in diesen Definitionen geradkettige und verzweigt-kettige Alkylgruppen und vollständig oder teilweise zyklisierte Alkylgruppen umfaßt und "Phenyl" substituierte Phenylgruppen umfaßt, welche bis zu 3 Substituenten enthalten, ausgewählt unter OH, C<sub>1</sub>-C<sub>4</sub>-Carboxy, Halogen, C<sub>1</sub>-C<sub>4</sub>-Alkyl, C<sub>1</sub>-C<sub>4</sub>-Alkoxy und C<sub>2</sub>-C<sub>4</sub>-Alkenyl.
3. Gemisch oder Kombination eines Produktes gemäß Anspruch 1 oder 2 mit einem pharmazeutisch oder tiermedizinisch verträglichen Träger zur Verwendung als ein Immunantwortmodifikator oder -regulator im Menschen oder in anderen Wirbeltierspezies.
4. Gemisch oder Kombination nach Anspruch 3, wobei der Träger ein subkutan verabreichbarer Träger ist.
5. Gemisch oder Kombination nach Anspruch 4, wobei der subkutan verabreichbare Träger ein lipophiler Träger ist.
6. Gemisch oder Kombination nach Anspruch 3, wobei der Träger ein lokal (einschließlich perkutan) verabreichbarer Träger ist.
7. Gemisch oder Kombination nach Anspruch 6, wobei der lokal verabreichbare Träger eine Lösung, eine Creme, eine Emulsion, eine Suspension, ein Gel, ein Puder, ein Spray, ein Pflaster oder ein Zäpfchen ist.
8. Gemisch oder Kombination nach Anspruch 3, wobei der Träger ein oral verabreichbarer Träger ist.
9. Verwendung von 5-Androsten-3 $\beta$ , 7 $\alpha$ , 17 $\beta$ -triol ( $\alpha$ AET) zur Herstellung eines Immunantwortmodifikators oder -regulators im Menschen oder in anderen Wirbeltierspezies.
10. Verwendung nach Anspruch 9, wobei die die Hydroxylgruppen schützenden Gruppen unter C<sub>1</sub>-C<sub>8</sub>-Alkyl, C<sub>2</sub>-C<sub>8</sub>-Alkenyl, Phenyl, Phenyl-(C<sub>1</sub>-C<sub>4</sub>)-Alkyl und -COR<sub>2</sub> ausgewählt sind, wobei R<sub>2</sub> H, C<sub>1</sub>-C<sub>8</sub>-Alkyl, C<sub>2</sub>-C<sub>8</sub>-Alkenyl, Phenyl oder Phenyl-(C<sub>1</sub>-C<sub>4</sub>)-Alkyl ist und wobei "Alkyl" in diesen Definitionen geradkettige und verzweigt-kettige Alkylgruppen und vollständig oder teilweise zyklisierte Alkylgruppen umfaßt und "Phenyl" substituierte Phenylgruppen umfaßt, welche bis zu 3 Substituenten enthalten, ausgewählt unter OH, C<sub>1</sub>-C<sub>4</sub>-Carboxy, Halogen, C<sub>1</sub>-C<sub>4</sub>-Alkyl, C<sub>1</sub>-C<sub>4</sub>-Alkoxy und C<sub>2</sub>-C<sub>4</sub>-Alkenyl.
11. Verwendung nach Anspruch 9 oder 10, wobei der Immunantwortmodifikator oder -regulator  $\alpha$ AET im Gemisch oder in Kombination mit einem pharmazeutisch oder tiermedizinisch verträglichen Träger enthält.
12. Verwendung nach Anspruch 11, wobei der Träger ein subkutan verabreichbarer Träger ist.
13. Verwendung nach Anspruch 12, wobei der subkutan verabreichbare Träger ein lipophiler Träger ist.

14. Verwendung nach Anspruch 11, wobei der Träger ein lokal (einschließlich perkutan) verabreichbarer Träger ist.
15. Verwendung nach Anspruch 14, wobei der lokal verabreichbare Träger eine Lösung, eine Creme, eine Emulsion, eine Suspension, ein Gel, ein Puder, ein Spray, ein Pflaster oder ein Zäpfchen ist.
- 5 16. Verwendung nach Anspruch 11, wobei der Träger ein oral verabreichbarer Träger ist.
17. Mit Hydroxylgruppenschutz versehenes Derivat von 5-Androsten-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED), wobei die Hydroxylgruppen durch Hydroxylgruppen schützende Gruppen geschützt sind, zur Verwendung als ein Immunantwortmodifikator oder -regulator im Menschen oder in anderen Wirbeltierspezies.
- 10 18. Mit Hydroxylgruppenschutz versehenes Derivat von  $\beta$ AED nach Anspruch 17, wobei die die Hydroxylgruppen schützenden Gruppen unter C<sub>1</sub>-C<sub>8</sub>-Alkyl, C<sub>2</sub>-C<sub>8</sub>-Alkenyl, Phenyl, Phenyl-(C<sub>1</sub>-C<sub>4</sub>)-Alkyl und -COR<sub>2</sub> ausgewählt sind, wobei R<sub>2</sub> H, C<sub>1</sub>-C<sub>8</sub>-Alkyl, C<sub>2</sub>-C<sub>8</sub>-Alkenyl, Phenyl oder Phenyl-(C<sub>1</sub>-C<sub>4</sub>)-Alkyl ist und wobei "Alkyl" in diesen Definitionen geradkettige und verzweigt-kettige Alkylgruppen und vollständig oder teilweise zyklisierte Alkylgruppen umfaßt und "Phenyl" substituierte Phenylgruppen umfaßt, welche bis zu 3 Substituenten enthalten, ausgewählt unter OH, C<sub>1</sub>-C<sub>4</sub>-Carboxy, Halogen, C<sub>1</sub>-C<sub>4</sub>-Alkyl, C<sub>1</sub>-C<sub>4</sub>-Alkoxy und C<sub>2</sub>-C<sub>4</sub>-Alkenyl.
- 15 19. Gemisch oder Kombination eines mit Hydroxylgruppenschutz versehenen Derivates von  $\beta$ AED nach Anspruch 17 oder 18 mit einem pharmazeutisch oder tiermedizinisch verträglichen Träger zur Verwendung als ein Immunantwortmodifikator oder -regulator im Menschen oder in anderen Wirbeltierspezies.
- 20 20. Gemisch oder Kombination nach Anspruch 19, wobei der Träger ein subkutan verabreichbarer Träger ist.
- 25 21. Gemisch oder Kombination nach Anspruch 20, wobei der subkutan verabreichbare Träger ein lipophiler Träger ist.
22. Gemisch oder Kombination nach Anspruch 19, wobei der Träger ein lokal (einschließlich perkutan) verabreichbarer Träger ist.
- 30 23. Gemisch oder Kombination nach Anspruch 22, wobei der lokal verabreichbare Träger eine Lösung, eine Creme, eine Emulsion, eine Suspension, ein Gel, ein Puder, ein Spray, ein Pflaster oder ein Zäpfchen ist.
24. Gemisch oder Kombination nach Anspruch 19, wobei der Träger ein oral verabreichbarer Träger ist.
- 35 25. Verfahren zur Verhinderung von Zelladhäsion, welches die Stufe umfaßt, bei der man die Zellen mit 5-Androsten-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED), 5-Androsten-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET) oder 5-Androsten-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET) in Kontakt bringt.
- 40 26. Verfahren zur Verstärkung von Lymphozytenproliferation in vitro, mit der Stufe, bei der man 5-Androsten-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) oder 5-Androsten-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET) in das Wachstumsmedium aufnimmt.
27. Verfahren nach Anspruch 26, angewendet auf die Verstärkung der Proliferation einer autogenen Lymphozytenpopulation aus einem Patienten vor der Rückverabreichung der verstärkten Lymphozytenpopulation an den Patienten.
- 45 28. Verfahren zur im wesentlichen stereospezifischen Herstellung von 5-Androsten-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol, welches die Stufe umfaßt, bei der man 3 $\beta$ ,17 $\beta$ -Diacetoxyandrost-5-en-7-on mit Aluminiumisopropoxid in Isopropanol reduziert.
- 50 29. Verfahren zur im wesentlichen stereospezifischen Herstellung von 5-Androsten-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol, welches die Stufe umfaßt, bei der man 3 $\beta$ ,17 $\beta$ -Diacetoxyandrost-5-en-7-on mit Triisobutylaluminium in Pentan unter Herstellung von 3 $\beta$ ,17 $\beta$ -Diacetoxy-7 $\beta$ -hydroxyandrost-5-en reduziert und anschließend die Acetoxygruppen entfernt.
30. Zusammensetzung, welche 5-Androsten-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) in einer Dosis von 0,2 bis 30 mg pro Tag in einem pharmazeutisch verträglichen Träger enthält, wobei der pharmazeutisch verträgliche Träger einen festen Träger umfaßt.
- 55 31. Zusammensetzung nach Anspruch 30, wobei der Träger ein Pflaster umfaßt.

32. Zusammensetzung nach Anspruch 30, wobei der Träger ein Kondom ist.
33. Verwendung der Zusammensetzung nach einem der Ansprüche 30 bis 32 für das Inkontaktbringen mit einer Scheidenwand.
- 5 34. Zusammensetzung nach einem der Ansprüche 30 bis 32 zur Verwendung als ein antivirales Mittel.
35. Kapsel, welche 5-Androsten-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) in einer Dosis von 0,2 bis 30 mg pro Tag in einem pharmazeutisch verträglichen Träger enthält.
- 10 36. Verwendung von 5-Androsten-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) zur Herstellung eines Arzneimittels zur Verhinderung oder Reduzierung einer Schädigung durch Strahlung im Menschen oder in anderen Wirbeltierspezies.
- 15 37. 5-Androsten-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET) zur Verwendung als ein aktives Mittel zur Verhinderung oder Reduzierung einer Schädigung durch Strahlung im Menschen oder in anderen Wirbeltierspezies.

# Revendications

- 20 1. 5-androstène-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET) ou un de ses dérivés à fonction hydroxy protégée, dans lequel les groupes hydroxy sont protégés par des groupes protecteurs hydrolysables, destiné à être utilisé comme modificateur ou régulateur de réponse immunitaire chez l'homme ou d'autres espèces de vertébrés.
- 25 2. Produit suivant la revendication 1, dans lequel les groupes protecteurs de la fonction hydroxy sont choisis entre des groupes alkyle en C<sub>1</sub> à C<sub>8</sub>, alcényle en C<sub>2</sub> à C<sub>8</sub>, phényle, phényl-(alkyle en C<sub>1</sub> à C<sub>4</sub>) et -COR<sub>2</sub>, dans lequel R<sub>2</sub> représente H, un groupe alkyle en C<sub>1</sub> à C<sub>8</sub>, alcényle en C<sub>2</sub> à C<sub>8</sub>, phényle ou phényl-(alkyle en C<sub>1</sub> à C<sub>4</sub>) ; le terme "alkyle" dans ces définitions désignant des groupes alkyle à chaîne droite et à chaîne ramifiée, et des groupes alkyle totalement ou partiellement cyclisés, et le terme "phényle" comprenant des groupes phényle substitués contenant jusqu'à 3 substituants choisis entre des substituants OH, carboxy en C<sub>1</sub> à C<sub>4</sub>, halogéno, alkyle en C<sub>1</sub> à C<sub>4</sub>, alkoxy en C<sub>1</sub> à C<sub>4</sub> et alcényle en C<sub>2</sub> à C<sub>4</sub>.
- 30 3. Mélange ou association d'un produit suivant la revendication 1 ou 2 avec un véhicule acceptable du point de vue pharmaceutique ou vétérinaire, destiné à être utilisé comme modificateur ou régulateur de réponse immunitaire chez l'homme ou d'autres espèces de vertébrés.
- 35 4. Mélange ou association suivant la revendication 3, dans lequel le véhicule est un véhicule pouvant être administré par voie sous-cutanée.
- 40 5. Mélange ou association suivant la revendication 4, dans lequel le véhicule pouvant être administré par voie sous-cutanée est un support lipophile.
6. Mélange ou association suivant la revendication 3, dans lequel le véhicule est un véhicule pouvant être administré par voie topique (y compris par voie percutanée).
- 45 7. Mélange ou association suivant la revendication 6, dans lequel le véhicule pouvant être administré par voie topique est une solution, une crème, une émulsion, une suspension, une gelée, une poudre, un liquide d'atomisation, un timbre ou un suppositoire.
- 50 8. Mélange ou association suivant la revendication 3, dans lequel le support est un véhicule pouvant être administré par voie orale.
9. Utilisation de 5-androstène-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET) pour la production d'un modificateur ou régulateur de réponse immunitaire chez l'homme ou d'autres espèces de vertébrés.
- 55 10. Utilisation suivant la revendication 9, dans laquelle les groupes protecteurs de la fonction hydroxy sont choisis entre des groupes alkyle en C<sub>1</sub> à C<sub>8</sub>, alcényle en C<sub>2</sub> à C<sub>8</sub>, phényle, phényl-(alkyle en C<sub>1</sub> à C<sub>4</sub>) et -COR<sub>2</sub>, dans lequel R<sub>2</sub> représente H, un groupe alkyle en C<sub>1</sub> à C<sub>8</sub>, alcényle en C<sub>2</sub> à C<sub>8</sub>, phényle ou phényl-(alkyle en C<sub>1</sub> à C<sub>4</sub>) ; le terme "alkyle" dans ces définitions désignant des groupes alkyle à chaîne droite et à chaîne ramifiée, et des

groupes alkyle totalement ou partiellement cyclisés, et le terme "phényle" comprenant des groupes phényle substitués contenant jusqu'à 3 substituants choisis entre des substituants OH, carboxy en C<sub>1</sub> à C<sub>4</sub>, halogéno, alkyle en C<sub>1</sub> à C<sub>4</sub>, alkoxy en C<sub>1</sub> à C<sub>4</sub> et alcényle en C<sub>2</sub> à C<sub>4</sub>.

- 5 11. Utilisation suivant la revendication 9 ou 10, dans laquelle le modificateur ou régulateur de réponse immunitaire comprend de l' $\alpha$ AET en mélange ou en association avec un véhicule acceptable du point de vue pharmaceutique ou vétérinaire.
- 10 12. Utilisation suivant la revendication 11, dans laquelle le véhicule est un support pouvant être administré par voie sous-cutanée.
13. Utilisation suivant la revendication 12, dans laquelle le véhicule pouvant être administré par voie sous-cutanée est un véhicule lipophile.
- 15 14. Utilisation suivant la revendication 11, dans laquelle le véhicule est un véhicule pouvant être administré par voie topique (y compris par voie percutanée).
- 20 15. Utilisation suivant la revendication 14, dans laquelle le véhicule pouvant être administré par voie topique est une solution, une crème, une émulsion, une suspension, une gelée, une poudre, un liquide d'atomisation, un timbre ou un suppositoire.
- 25 16. Utilisation suivant la revendication 11, dans laquelle le support est un véhicule pouvant être administré par voie orale.
- 30 17. Dérivé à fonction hydroxy protégée de 5-androstène-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED), dans lequel les groupes hydroxy sont protégés par des groupes protecteurs de la fonction hydroxy, destiné à être utilisé comme modificateur ou régulateur de réponse immunitaire chez l'homme ou d'autres espèces de vertébrés.
- 35 18. Dérivé à fonction hydroxy protégée de  $\beta$ AED suivant la revendication 17, dans lequel les groupes protecteurs de la fonction hydroxy sont choisis entre des groupes alkyle en C<sub>1</sub> à C<sub>8</sub>, alcényle en C<sub>2</sub> à C<sub>8</sub>, phényle, phényl-(alkyle en C<sub>1</sub> à C<sub>4</sub>) et -COR<sub>2</sub>, dans lequel R<sub>2</sub> représente H, un groupe alkyle en C<sub>1</sub> à C<sub>8</sub>, alcényle en C<sub>2</sub> à C<sub>8</sub>, phényle ou phényl-(alkyle en C<sub>1</sub> à C<sub>4</sub>) ; le terme "alkyle" dans ces définitions désignant des groupes alkyle à chaîne droite et à chaîne ramifiée, et des groupes alkyle totalement ou partiellement cyclisés, et le terme "phényle" comprenant des groupes phényle substitués contenant jusqu'à 3 substituants choisis entre des substituants OH, carboxy en C<sub>1</sub> à C<sub>4</sub>, halogéno, alkyle en C<sub>1</sub> à C<sub>4</sub>, alkoxy en C<sub>1</sub> à C<sub>4</sub> et alcényle en C<sub>2</sub> à C<sub>4</sub>.
- 40 19. Mélange ou association d'un dérivé, à fonction hydroxy protégée, de  $\beta$ AED suivant la revendication 17 ou 18 avec un véhicule acceptable du point de vue pharmaceutique ou vétérinaire, destiné à être utilisé comme modificateur ou régulateur de réponse immunitaire chez l'homme ou d'autres espèces de vertébrés.
- 45 20. Mélange ou association suivant la revendication 19, dans lequel le support est un véhicule pouvant être administré par voie sous-cutanée.
- 50 21. Mélange ou association suivant la revendication 20, dans lequel le véhicule pouvant être administré par voie sous-cutanée est un véhicule lipophile.
- 55 22. Mélange ou association suivant la revendication 19, dans lequel le véhicule est un véhicule pouvant être administré par voie topique (y compris par voie percutanée).
23. Mélange ou association suivant la revendication 22, dans lequel le véhicule pouvant être administré par voie topique est une solution, une crème, une émulsion, une suspension, une gelée, une poudre, un liquide d'atomisation, un timbre ou un suppositoire.
24. Mélange ou association suivant la revendication 19, dans lequel le support est un véhicule pouvant être administré par voie orale.
25. Méthode pour la prévention d'une adhésion cellulaire, qui comprend la mise en contact des cellules avec du 5-androstène-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED), du 5-androstène-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET) ou du 5-androstène-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET).

26. Méthode pour accroître la prolifération des lymphocytes *in vitro*, qui comprend l'incorporation au milieu de croissance de 5-androstène-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) ou du 5-androstène-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol ( $\beta$ AET).
- 5 27. Méthode suivant la revendication 26, appliquée à l'accroissement de la prolifération d'une population de lymphocytes autogènes provenant d'un patient avant la réadministration de la population de lymphocytes accrue au patient.
- 10 28. Méthode pour la production substantiellement stéréospécifique de 5-androstène-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol, qui comprend l'étape consistant à réduire la 3 $\beta$ ,17 $\beta$ -diacétoxyandrost-5-ène-7-one avec de l'isopropylate d'aluminium dans de l'isopropanol.
- 15 29. Méthode pour la production substantiellement stéréospécifique de 5-androstène-3 $\beta$ ,7 $\beta$ ,17 $\beta$ -triol, qui comprend l'étape de réduction de la 3 $\beta$ ,17 $\beta$ -diacétoxyandrost-5-ène-7-one avec du triisobutylaluminium dans le pentane pour produire du 3 $\beta$ ,17 $\beta$ -diacétoxy-7 $\beta$ -hydroxyandrost-5-ène et ensuite l'élimination des groupes acétoxy.
30. Composition comprenant du 5-androstène-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) à une dose de 0,2 à 30 mg par jour dans un véhicule pharmaceutiquement acceptable, le véhicule pharmaceutiquement acceptable comprenant un support solide.
- 20 31. Composition suivant la revendication 30, dans laquelle le support comprend un timbre.
32. Composition suivant la revendication 30, dans laquelle le support est un préservatif.
33. Utilisation de la composition suivant l'une quelconque des revendications 30 à 32 pour la mise en contact avec une paroi vaginale.
- 25 34. Composition suivant l'une quelconque des revendications 30 à 32, destinée à être utilisée comme agent antiviral.
35. Capsule comprenant du 5-androstène-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) à une dose de 0,2 à 30 mg par jour dans un véhicule pharmaceutiquement acceptable.
- 30 36. Utilisation de 5-androstène-3 $\beta$ ,17 $\beta$ -diol ( $\beta$ AED) pour la production d'un médicament destiné à la prévention ou à la réduction de l'altération par irradiation chez l'homme ou d'autres espèces de vertébrés.
- 35 37. 5-androstène-3 $\beta$ ,7 $\alpha$ ,17 $\beta$ -triol ( $\alpha$ AET) destiné à être utilisé comme agent actif pour la prévention ou la réduction de l'altération par irradiation chez l'homme ou d'autres espèces de vertébrés.